

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

317

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 31/55, 47/00, 47/10, 47/12</b>		<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 95/29678</b>
			<b>(43) International Publication Date:</b> 9 November 1995 (09.11.95)
<b>(21) International Application Number:</b> PCT/EP95/01519			<b>(81) Designated States:</b> AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, IS, JP, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).
<b>(22) International Filing Date:</b> 21 April 1995 (21.04.95)			
<b>(30) Priority Data:</b> 234,216 28 April 1994 (28.04.94) US 234,214 28 April 1994 (28.04.94) US			
<b>(71) Applicant (for all designated States except US):</b> F.HOFFMANN-LA ROCHE AG [CH/CH]; Grenzacherstrasse 124, CH-4002 Basle (CH).			
<b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> BEHL, Charanjit, Rai [US/US]; 27 Cottage Place, Nutley, NJ 07110 (US). HOFFMANN, Peter [DE/DE]; Ruthardweg 17, D-79576 Weil-Haltingen (DE).			
<b>(74) Agent:</b> GROSSNER, Lutz; Grenzacherstrasse 124, CH-4002 Basle (CH).			<b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> PHARMACEUTICAL COMPOSITION FOR TRANSDERMAL DELIVERY			
<b>(57) Abstract</b>  A pharmaceutical composition for transdermal delivery comprising an effective amount of an active ingredient selected from a benzodiazepine and a benzodiazepine antagonist; and either ethanol; caprylic acid; and oleic acid; or isopropanol, propylene glycol, oleic acid, and water. Additionally, the composition may contain silicon fluid, benzyl alcohol, transcutol or dimethyl sulfoxide.			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgystan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Larvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

## Pharmaceutical composition for transdermal delivery

Benzodiazepines are used as sedative hypnotics, in the treatment of anxiety disorders and in the treatment of seizures.

- 5        Benzodiazepine antagonists, such as, flumazenil, are used for a complete or partial reversal of the sedative effects of benzodiazepines and for the management of benzodiazepine overdose.

10       Benzodiazepines and benzodiazepine antagonists, are administered either via gastrointestinal tract or parenterally. Alternatively, a transdermal route of drug delivery can be used. Generally, the most critical problem in this route is the lack of adequate absorption of drugs through the skin.

- 15       A pharmaceutical composition for transdermal delivery comprising an effective amount of an active ingredient selected from a benzodiazepine and a benzodiazepine antagonist, and either a) caprylic acid, ethanol, and oleic acid; or b) isopropanol, propylene glycol, oleic acid, and water.

20

As used herein, the term benzodiazepine means any active pharmaceutical compound in the benzodiazepine family, such as, diazepam, chlordiazepoxide, fluazepam, lorazepam and clonazepam, preferably clonazepam.

25

As used herein the term benzodiazepine antagonist means any compound antagonistic to benzodiazepines, such as, preferably flumazenil.

30

In one aspect, the present invention relates to a pharmaceutical composition for transdermal delivery comprising an effective amount

of an active ingredient selected from a benzodiazepine and benzodiazepine antagonist; ethanol; caprylic acid; and oleic acid with or without an inert carrier.

5        Ethanol is present preferably in the range of from about 10 to about 95 percent by weight of the compositions. In a particularly preferred embodiment, ethanol is present in the composition in the range of from about 24 to about 90 percent by weight of such compositions.

10

Preferably, caprylic acid is present in such compositions in the range of from about 1 to about 10 percent by weight of the composition, particularly preferred at about 3 percent.

15        In the compositions according to that aspect of the invention, preferably, oleic acid is present in such compositions in the range of from about 1 to about 10 percent by weight of the composition, particularly preferred at about 3 percent.

20        When the active ingredient is a benzodiazepine antagonist, such as flumazenil, preferably, ethanol is present in such compositions in an amount of from about 10 to about 95 percent by weight of the composition; particularly preferred in an amount of from about 50 to about 70 percent; caprylic acid is present in the composition in an  
25        amount of from about 1 to about 10 percent by weight of the composition, particularly preferred in an amount of from about 3 to 5 percent; and oleic acid is present in such compositions in an amount of from about 1 to about 10 percent by weight of the composition, particularly preferred in an amount of from about 3 to 5 percent.

30

When the active ingredient is a benzodiazepine, such as clonazepam, preferably, ethanol is present in the composition in an amount of from about 10 to about 95 percent by weight of the composition, particularly preferred in an amount of from about 50 to  
35        90 percent; caprylic acid is present in the composition in an amount of

## 3

from about 1 to about 10 percent by weight of the composition, particularly preferred at about 3 percent; and oleic acid is present in the composition in an amount of from about 1 to about 10 percent by weight of the composition, particularly preferred at about 3 percent.

5 The compositions of the above described aspect of the invention may contain additional enhancing materials such as, for example, silicon fluid such as Silicon Dow® 556 (polyphenyl methyl siloxane), preferably in the range of from about 15 to about 25 percent by  
10 weight of the composition, particularly preferred at about 20 percent; dimethylsulfoxide, preferably in the range of from about 1 to about 20 percent by weight of the composition, particularly preferred at about 2 percent; acetone, preferably in the range of from about 15 to about 25 percent by weight of the composition, particularly preferred  
15 at about 20 percent; caprylic/capric triglyceride such as Miglyol® 840 (propylene glycol diesters of saturated vegetable fatty acids of the chain lengths C8-C10, particularly 2% max caproic acid (C6:0), 65-80% caprylic acid (C8:0), 15-30% capric acid (C10:0), and 3% max. linoleic acid (C18:2) Dynamit Nobel), preferably in the range of from about 25  
20 to about 40 percent by weight of the composition, particularly preferred at about 36 percent; transcitol (diethylene glycol monoethyl ether from Gattefosse) preferably in the range of from about 15 to about 30 percent by weight of the composition, particularly preferred at about 20 percent; and benzyl alcohol,  
25 preferably in the range of from about 5 to about 15 percent by weight of the composition, particularly preferred at about 10 percent.

In another aspect, the present invention relates to a pharmaceutical composition for transdermal delivery comprising an effective  
30 amount of an active ingredient selected from a benzodiazepine and benzodiazepine antagonist; isopropanol; propylene glycol; oleic acid; and water with or without an inert carrier.

Preferably, isopropanol is present in such compositions in the  
35 range of from about 10 to about 95 percent by weight of the

composition. In a particularly preferred embodiment, isopropanol is present in the composition in an amount of about 20% by weight of the composition.

5            Preferably, propylene glycol is present in these compositions in the range of from about 30 to about 50 percent by weight of the composition, particularly preferred in the range of from about 35 to about 45 percent by weight.

10           Preferably, oleic acid is present in these compositions in the range of from about 1 to about 10 percent by weight of the composition, particularly preferred is about 5 percent by weight.

15           Preferably, water is present in these compositions in the range of from about 10 to about 30 percent by weight of the composition, particularly preferred in the range of from about 20 to about 25 percent by weight.

20           When the active ingredient is a benzodiazepine antagonist such as flumazenil, preferably isopropanol is present in these compositions in an amount of about 20 percent by weight of the composition; propylene glycol is present in an amount of from about 38 to about 47 percent by weight of the composition; oleic acid is present in an amount of about 5 percent by weight of the composition; and water is  
25           present in an amount of about 20 percent by weight of the composition.

30           The composition of the aforesaid aspect of the invention may contain additional enhancing materials such as, for example, Diacetin (glycerol diacetate from Davos Chemical), preferably in the range of from 5 to 15 percent by weight of the composition, particularly preferred in about 10 percent; Cetiol B® (dibutyl adipate from Henkel Co.), preferably in the range of from 1 to 10 percent by weight of the composition, particularly preferred in about 5 percent; caprylic acid,  
35           preferably in the range of from 1 to 10 percent of the composition,

- 5 -

particularly preferred in about 5 percent; silicon fluid such as, Silicon Dow® 556 (polyphenyl methyl siloxane), preferably in the range of from 5 to 15 percent by weight of the composition, particularly preferred in about 10 percent; caprylic/capric triglyceride, such as Miglyol® 840 (propylene glycol diesters of saturated vegetable fatty acids of the chain lengths C8-C10, particularly 2% max caproic acid (C6:0), 65-80% caprylic acid (C8:0), 15-30% capric acid (C10:0), and 3% max. linoleic acid (C18:2) Dynamit Nobel) preferably in the range of from 5 to 15 percent by weight of the composition, particularly preferred in about 10 percent; transcitol (diethylene glycol monoethyl ether from Gattefosse), preferably in the range of from 5 to 15 percent by weight of the composition, particularly preferred in about 10 percent.

Pharmaceutical compositions in accordance with this invention can be formulated to additionally contain conventional additives or supplementary ingredients in the usual amounts for such materials. The composition can be in the form of a gel, as well as, in the form of a solution, preferably a thickened solution. By way of illustration such additives or supplements include the following.

Gelling agents which can be used include, for example, hydroxy methyl cellulose, preferably in the range of from 1 to 4 percent by weight of the composition; tragacanth preferably in the range of from 2 to 5 percent by weight of the composition; sodium alginate, preferably in the range of from 2 to 10 percent by weight of the composition; gelatin, preferably in the range of from 2 to 15 percent by weight of the composition; methylcellulose, preferably in the range of from 2 to 4 percent by weight of the composition; sodium carboxymethylcellulose, preferably in the range of from 2 to 5 percent by weight of the composition; and polyvinyl alcohols, preferably in the range of from 10 to 20 percent by weight of the composition. A particularly preferred gelling agent is Klucel®.

Klucel HF is a hydroxypropyl cellulose (Hercules Inc.) with a molecular weight in the 1,000,000 range and moisture content of 17% for 1,500-2,500. Hydroxypropyl cellulose is preferably present in the composition in the range of from 1.0 to 5.0 percent by weight, particularly preferred in the range of from 1.0 to 4.0 percent by weight. Generally, enough Klucel is added to provide a reasonably good gel-consistency to the product.

The preservatives which can be used in the invention include, for example, parabens, preferably at about 0.2%; benzoic acid, preferably at about 0.2%; and, chlorocresol, preferably at about 0.1%.

If needed, antioxidants can be used in the gel formulations to improve the stability of the drug. These antioxidants include, for example, ascorbyl palmitate, butylated hydroxyanisole, butylated hydroxytoluene, potassium sorbate, sodium bisulfate, sorbic acid, propyl gallate and sodium metabisulfite.

Preferably, the pharmaceutical composition of the invention are administered to a host in need of such treatment in a transdermal patch of a reservoir type.

Adhesives used in making transdermal patches for use with the invention include, for example, preferably poly-isobutylene, silicone based adhesives and acrylic polymers. The adhesive polymers can be mixed with other excipients such as mineral oil to make them more suitable for a given purpose.

The backing membrane of a transdermal patch constitutes the upper part (exposed to the environment) of a transdermal patch and is made of materials such as, for example, preferably polyester films, ethyl vinyl acetate, polypropylene, polyethylene and polyvinyl-chloride.



A rate controlling membrane of a transdermal patch is placed in contact with the pharmaceutical composition of the invention and its other side is in contact with the skin of a host. The rate controlling membrane is made of materials such as, for example, preferably,  
5 dimethylpolysiloxane, polyacetate, polyurethane and ethylene-vinyl acetate copolymer and polypropylene.

At the bottom of a transdermal patch, a protective liner is placed in contact with the adhesive layer. This liner protects against  
10 the drug release from the formulation reservoir until the liner is peeled off the patch and applied on the skin surface of the host. Such liners are made of materials including preferably polyethylene terephthalate film, polyester membrane and polycarbonate film.

15 Alternatively, one can make transdermal patches which are called monolithic or adhesive type patches. In this case, the drug is dispersed either in a suitable adhesive or in a suitable non-adhesive polymer and then the mixture is layered onto a membrane. A protective membrane is placed on the adhesive.

20

In vivo tests were utilized to evaluate the absorption of benzodiazepines and benzodiazepine antagonists administered in accordance with this invention.

25

### Methods

General Procedure: Hairless guinea pigs (HGP) were anesthetized by using Ketamin-HCl and promazine. The side sites of the animals were cleaned with water. Zero time blood samples were withdrawn from  
30 the ocular site. The transdermal drug delivery systems were placed on the skin, two per animal providing a total area of 9.0 to 10.0 sq. cm., precisely measured. The animals were allowed to come out of anesthesia in between blood samples. Blood samples were withdrawn at 1.0, 2.0, 3.0, 4.0, and 6.0 hours. The blood was allowed to clot and  
35 then centrifuged to obtain serum. The drug concentration was

determined by using an HPLC method. After the last sample point, the transdermal drug delivery system was removed from the animal's skin and the site was examined for any "obvious" signs of irritation/reddening.

5

Serum Collection: The animals were bled from the eye into Microtainer serum separator tubes (Becton Dickinson, 5960). The blood (0.6 mL) was centrifuged at 4,000 rpm for 15 minutes (4,400 g) on a Beckman J-6M centrifuge with a JS-4.2 rotor. Serum was  
10 separated and frozen until the HPLC analysis. Before sample preparation, the serum was thawed and centrifuged again.

Sample Preparation: Two hundred and fifty microliters of serum were mixed with 250 mcL of water and 25 mcL of an internal  
15 standard, flunitrazepam 1 mcg/mL in methanol, were added. The sample was purified on a solid phase mini column, Adsorbex RP-18 (100 mg; EM Science) using the sample preparation unit Adsorbex SPU). The columns were treated before with 2 mL of methanol and washed with 4 mL of water. Samples were applied and the columns  
20 were washed with 4 mL of water. The columns were dried under vacuum (5" Hg) and eluted with two portions of 125 mcL of acetonitrile:water (1:1).

HPLC Conditions: Samples were analyzed on a Waters HPLC system  
25 using Waters 600E controller, Waters 712 WISP automatic sample injector and Applied Biosystems 785A programmable absorbance detector.

---

Column:	Waters Nova Pak C18, 75 X 3.9 mm
Flow rate:	2.0 mL/min
5 Mobile Phase:	25% acetonitrile in water (v/v)
Wavelength:	310 nm
Data collection:	2 points/sec, 1 V/AU, A/D = 0.1, rise time=1 sec
Injection volume:	100 mcL
10 Run time:	15 min/sample

---

The retention times of the internal standard, flunitrazepam, and clonazepam or flumazenil were 5.5 and 4.5 minutes, respectively. The HPLC system is connected to a computer where a program was used to determine the area under the curve of the drug and the internal standard.

Lack Of Interference: The chromatogram of the HGP serum shows no peak at the retention time of clonazepam indicating an interference free detection of the drug.

Sensitivity And Linearity Of Response: A standard curve was made by adding clonazepam or flumazenil and the internal standard to HGP serum. A linear relationship was observed between the observed response and concentration of clonazepam in the range of 5 to 500 ng/mL. The recovery of the drug in these experiments was  $75 \pm 15\%$ , and was corrected using the internal standard. Apparent limit of quantification was found to be about 5 ng/mL of clonazepam or flumazenil in the HGP serum.

Data Analysis: The HPLC data were computed in terms of drug concentration per unit volume of the serum and were plotted as a function of time. In such experiments, the blood levels are expected to rise to a maximum and then decline due to a decrease in the chemical

potential of the drug in the patch. No rate controlling membrane was placed at the bottom of the contemporary transdermal delivery dosage system.

5

ResultsTable I

10	<u>Formulation<sup>a</sup></u>	<u>Max Blood Level Observed in HGP (ng/ml)</u>
	Example 1	130
	Example 2	90
	Example 3	80
15	Example 4	300
	Example 5	37
	Example 6	525
	Example 7	470
	Example 8	225
20	Example 9	273
	Control A <sup>b</sup>	33
	Control B <sup>c</sup>	15

a Dose was 12 mg per animal applied to an area of 9 cm sq.

25 b Control A contained 11 mg of clonazepam in a formulation comprised of 97% ethanol and 3% Klucel HF applied to an area of 9.8 cm<sup>2</sup>

c Control B contained 12 mg of flumazenil in a formulation comprised of 97% ethanol and 3% Klucel HF applied to an area of 30 5.0 cm<sup>2</sup>.

ResultsTable II

5	<u>Formulation<sup>a</sup></u>	<u>Max Blood Level Observed in HPG (ng/ml)</u>
	Example 10	762
	Example 11	733
10	Example 12	753
	Example 13	200
	Example 14	377
	Example 15	249
	Example 16	530
15	Control A <sup>b</sup>	15

<sup>a</sup> Drug concentration was 10 mg/Gm;

Dose was 12 mg per animal applied to an area of 9 cm sq.

<sup>b</sup> Control A contained 12 mg of flumazenil in a formulation comprised of 97% ethanol and 3% Klucel HF

By way of illustration, some suitable pharmaceutical compositions in accordance with this invention are set forth below. While clonazepam and flumazenil, the preferred benzodiazepine and benzodiazepine antagonist for the non-aqueous compositions and flumazenil, the preferred benzodiazepine antagonist for the aqueous compositions of this invention, are used to illustrate the compositions, it should be understood that other benzodiazepine and benzodiazepine antagonists may be substituted in appropriate amounts.

EXAMPLE 1

35	Clonazepam	0.010 Gm
	Ethanol	0.900 Gm
	Caprylic Acid	0.030 Gm
	Oleic Acid	0.030 Gm
	<u>Klucel HF</u>	<u>0.030 Gm</u>
	Total	1.000 Gm

EXAMPLE 2

	Clonazepam	0.010 Gm
	Ethanol	0.702 Gm
5	Silicon Fluid	0.198 Gm
	Caprylic Acid	0.030 Gm
	Oleic Acid	0.030 Gm
	<u>Klucel HF</u>	<u>0.030 Gm</u>
	Total	1.000 Gm

10

EXAMPLE 3

	Clonazepam	0.010 Gm
	Ethanol	0.800 Gm
15	Benzyl Alcohol	0.100 Gm
	Caprylic Acid	0.030 Gm
	Oleic Acid	0.030 Gm
	<u>Klucel HF</u>	<u>0.030 Gm</u>
	Total	1.000 Gm

20

EXAMPLE 4

	Clonazepam	0.010 Gm
	Ethanol	0.600 Gm
25	Silicon Fluid	0.200 Gm
	Benzyl Alcohol	0.100 Gm
	Caprylic Acid	0.030 Gm
	Oleic Acid	0.030 Gm
	<u>Klucel HF</u>	<u>0.030 Gm</u>
30	Total	1.000 Gm

EXAMPLE 5

	Clonazepam	0.010 Gm
	Ethanol	0.510 Gm
5	Transcutol	0.200 Gm
	Silicon Fluid	0.200 Gm
	Caprylic Acid	0.030 Gm
	Oleic Acid	0.030 Gm
	<u>Klucel HF</u>	<u>0.020 Gm</u>
10	Total	1.000 Gm

EXAMPLE 6

	Flumazenil	0.010 Gm
15	Ethanol	0.710 Gm
	Silicon Dow 556	0.20 Gm
	Caprylic Acid	0.03 Gm
	Oleic Acid	0.03 Gm
	<u>Klucel HF</u>	<u>0.02 Gm</u>
20	Total	1.000 Gm

EXAMPLE 7

	Flumazenil	0.010 Gm
25	Ethanol	0.690 Gm
	Silicon Dow 556	0.200 Gm
	Dimethyl Sulfoxide	0.020 Gm
	Oleic Acid	0.030 Gm
	Caprylic Acid	0.030 Gm
30	<u>Klucel HF</u>	<u>0.020 Gm</u>
	Total	1.000 Gm

EXAMPLE 8

	Flumazenil	0.010 Gm
	Ethanol	0.510 Gm
5	Acetone	0.200 Gm
	Silicon DOW 556	0.200 Gm
	Caprylic Acid	0.030 Gm
	Oleic Acid	0.030 Gm
	<u>Klucel HF</u>	<u>0.020 Gm</u>
10	Total	1.000 Gm

EXAMPLE 9

	Flumazenil	0.010 Gm
15	Ethanol	0.24 Gm
	Transcutol	0.250 Gm
	Miglyol 840	0.360 Gm
	Caprylic Acid	0.050 Gm
	Oleic Acid	0.050 Gm
20	<u>Klucel HF</u>	<u>0.040 Gm</u>
	Total	1.000 Gm

EXAMPLE 10

25	Flumazenil	0.010 Gm
	Isopropanol	0.205 Gm
	Propylene Glycol	0.410 Gm
	Oleic Acid	0.050 Gm
	Water	0.205 Gm
30	Klucel HF	0.010 Gm
	<u>Diacetin</u>	<u>0.110 Gm</u>
	Total	1.000 Gm



EXAMPLE 11

	Flumazenil	0.010 Gm
	Isopropanol	0.200 Gm
5	Propylene Glycol	0.380 Gm
	Water	0.200 Gm
	Oleic Acid	0.050 Gm
	Klucel HF	0.010 Gm
	Diacetin	0.100 Gm
10	<u>Cetiol B</u>	<u>0.050 Gm</u>
	Total	1.000 Gm

EXAMPLE 12

15	Flumazenil	0.010 Gm
	Isopropanol	0.220 Gm
	Propylene Glycol	0.440 Gm
	Water	0.220 Gm
	Caprylic Acid	0.050 Gm
20	Oleic Acid	0.050 Gm
	<u>Klucel HF</u>	<u>0.010 Gm</u>
	Total	1.000 Gm

EXAMPLE 13

25	Flumazenil	0.010 Gm
	Isopropanol	0.198 Gm
	Propylene Glycol	0.426 Gm
	Water	0.198 Gm
30	Silicon Dow 556	0.099 Gm
	Oleic Acid	0.049 Gm
	<u>Klucel HF</u>	<u>0.020 Gm</u>
	Total	1.000 Gm

EXAMPLE 14

	Flumazenil	0.010 Gm
	Isopropanol	0.228 Gm
5	Propylene Glycol	0.465 Gm
	Water	0.228 Gm
	Oleic Acid	0.049 Gm
	<u>Klucel HF</u>	<u>0.020 Gm</u>
	Total	1.000 Gm

10

EXAMPLE 15

	Flumazenil	0.010 Gm
	Isopropanol	0.198 Gm
15	Propylene Glycol	0.426 Gm
	Water	0.198 Gm
	Miglyol 840	0.099 Gm
	Oleic Acid	0.049 Gm
	<u>Klucel HF</u>	<u>0.020 Gm</u>
20	Total	1.000 Gm

EXAMPLE 16

	Flumazenil	0.010 Gm
25	Isopropanol	0.198 Gm
	Propylene Glycol	0.426 Gm
	Water	0.198 Gm
	Oleic Acid	0.049 Gm
	Transcutol	0.099 Gm
30	<u>Klucel HF</u>	<u>0.020 Gm</u>
	Total	1.000 Gm

The various ingredients of the formulations were mixed together in a glass apparatus. The drug was dissolved in this mixture. The gelling agent was added to this solution and the contents were mixed by using shear provided by a magnetic stirrer.

CLAIMS

1. A pharmaceutical composition for transdermal delivery comprising an effective amount of an active ingredient selected from a benzodiazepine and a benzodiazepine antagonist, and either a) caprylic acid, ethanol, and oleic acid; or b) isopropanol, propylene glycol, oleic acid, and water.
2. The composition of claim 1, wherein the benzodiazepine is clonazepam.
3. The composition of claim 1, wherein the benzodiazepine antagonist is flumazenil.
4. The composition of any one of claims 1-3, wherein ethanol is present in an amount of from about 10 to about 95 percent by weight of the composition; caprylic acid is present in an amount of from about 1 to about 10 percent by weight of the composition and oleic acid is present in an amount of from about 1 to about 10 percent by weight of the composition.
5. The composition of any one of claims 1-3, wherein isopropanol is present in an amount of from about 10 to about 95 percent by weight of the composition, propylene glycol is present in an amount of from about 30 to about 50 percent by weight of the composition, oleic acid is present in an amount of from about 1 to about 10 percent by weight of the composition, and water is present in an amount from about 10 to about 30 percent by weight of the composition.
6. The composition of claim 4 or 5, further comprising hydroxypropyl cellulose in an amount of from about 1 to about 4 percent by weight of the composition.
7. The composition of claim 4 or 5, further comprising silicon fluid.

8. The composition of claim 4, further comprising benzyl alcohol.
9. The composition of claim 4, further comprising dimethyl sulfoxide.
- 5 10. The composition of claim 4, further comprising acetone.
11. The composition of claim 4, further comprising diethyl glycol monoethyl ether.
- 10 12. The composition of claim 4 or 5, further comprising caprylic/capric triglyceride.
13. The composition of claim 5, further comprising glycerol diacetate.
- 15 14. The composition of claim 13, further comprising dibutyl adipate.
- 15 15. The invention as described hereinbefore, especially with reference to the Examples.

\*\*\*

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 95/01519

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K31/55 A61K47/00 A61K47/10 A61K47/12

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP,A,0 159 167 (TAKEDA CHEMICAL INDUSTRIES) 23 October 1985 see claims 1,9 see page 4, line 6 - page 6, line 3 see page 7, line 34 - page 9, line 23 see page 12, line 1 - page 14, line 20 ---	1-15
A	CH,A,634 749 (KALI-CHEMIE PHARMA) 28 February 1983 see claims 1-3,9,10 see page 3, right column, line 14 - line 31 -----	1-15



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

25 July 1995

Date of mailing of the international search report

27.08.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,

Authorized officer

Ventura Amat. A

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 95/01519

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0159167	23-10-85	JP-A- 61186311	20-08-86
		JP-A- 60214730	28-10-85
		CA-A- 1249968	14-02-89
		US-A- 4952560	28-08-90
<hr/>			
CH-A-634749	28-02-83	US-A- 4722941	02-02-88
<hr/>			